



RESEARCH ARTICLE

Prognostic value of plasminogen activator inhibitor-1 in biomarker exploration using multiplex immunoassay in patients with metastatic renal cell carcinoma treated with axitinib

Naoko Honma¹ | Takamitsu Inoue^{1,2}  | Norihiko Tsuchiya^{1,3} | Atsushi Koizumi¹ | Ryohei Yamamoto¹ | Taketoshi Nara¹ | Sohei Kanda¹ | Mingguo Huang¹ | Kazuyuki Numakura¹  | Mitsuru Saito¹ | Shintaro Narita^{1,2} | Shigeru Satoh⁴ | Tomonori Habuchi^{1,2}

¹Department of Urology, Akita University Graduate School of Medicine, Akita, Japan

²AMED-CREST, Japan Science and Technology Agency, Tokyo, Japan

³Department of Urology, Yamagata University Faculty of Medicine, Yamagata, Japan

⁴Center for Kidney Disease and Transplantation, Akita University Hospital, Akita, Japan

Correspondence

Takamitsu Inoue, Department of Renal and Urological Surgery, International University of Health and Welfare Narita Hospital, 852 Hatakeda, Narita, Chiba 286-0124, Japan. Email: takmitz@gmail.com; takmitz@iuhw.ac.jp

Funding information

Japanese Society for the Promotion of Science, Grant/Award Numbers: 16H02679, 23590168, 25293332

Abstract

Background and Aims: Vascular endothelial growth factor-directed therapies play a significant role in patients with metastatic renal cell carcinoma (mRCC). Biomarkers for predicting treatment efficacy and resistance are required to develop personalized medicine. We evaluated multiple serum cytokine levels in patients with mRCC treated with axitinib to explore predictive biomarkers.

Methods: From September 2012 to October 2015, serum samples were collected from 44 patients with mRCC before treatment and 4 weeks after axitinib initiation. Bio-Plex Pro Human Cancer Biomarker Panels 1 and 2 were used to measure levels of 34 serum biomarkers related to angiogenesis and cell proliferation.

Results: Patients with partial response or stable disease had significantly decreased serum plasminogen activator inhibitor-1 (PAI-1) level from pre-treatment to 4 weeks after axitinib initiation compared with those with progressive disease ($P = .022$). The median progression-free survival (PFS) and median overall survival (OS) in patients with increased serum PAI-1 level from pre-treatment to 4 weeks after axitinib initiation were significantly shorter than those with decreased serum PAI-1 level ($P = .027$ and $P = .026$, respectively). Increased serum PAI-1 level from pre-treatment to 4 weeks after axitinib initiation was an independent prognostic marker for shorter PFS and OS in multivariate analyses ($P = .015$ and $P = .032$, respectively). The immunohistochemical staining intensity of PAI-1 in tumor specimens was significantly associated with Fuhrman grade and presence of distant metastasis ($P = .026$ and $P = .010$, respectively).

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2020 The Authors. *Health Science Reports* published by Wiley Periodicals LLC.

Conclusions: The initial change in serum PAI-1 level in the early stage of axitinib treatment could be a useful prognostic biomarker in patients with mRCC.

KEYWORDS

metastatic renal cell carcinoma, molecular-targeted therapy, plasminogen activator inhibitor-1, serum biomarker

1 | INTRODUCTION

In 2017, the age-adjusted incidence and mortality rates of renal cell carcinoma (RCC) in Japanese men were 11.5 and 2.8 per 100 000 person-years, respectively.¹ Distant metastasis is observed in approximately 20% to 30% of patients with RCC at the time of initial diagnosis.² Although current first-line treatment for patients with metastatic RCC (mRCC) is either an immune-checkpoint inhibitor (ICI) or vascular endothelial growth factor (VEGF)-directed multitargeted tyrosine kinase inhibitors (TKIs),³ TKIs improved overall survival (OS) in patients with mRCC with a median value of 8.5 to 14.4 months from 2002 to 2008.⁴ Although the treatment paradigm for mRCC is currently shifting from TKIs to ICIs with or without concurrent use of TKIs, personalized biomarker-guided sequential or combination therapies for predicting the efficacy and adverse effects of TKIs are still strongly required for patients with mRCC.³

For appropriate use of TKIs in individual patients, useful biomarkers which can be measured during treatment to predict treatment effect, resistance, and prognosis are strongly required. As strategies to predict the treatment effect and prognosis during treatment, serum TKI level can be measured.⁵ Pre-treatment evaluation of genetic polymorphisms of drug-metabolizing enzymes and transporters can predict the serum TKI level.⁵ In addition, serum VEGF-C, sVEGFR-2, and sVEGFR-3 levels,⁶⁻⁸ and the number of endothelial cells in circulating blood⁹ have been reported to be biomarkers that correlate with treatment effect and prognosis. However, other potential biomarkers relevant to personalized therapy including TKIs and immunotherapies have not been investigated.

Axitinib is a TKI selective for VEGFR-1, -2, and -3. Patients with mRCC treated with axitinib as second-line therapy had a significantly longer progression-free survival (PFS) than those treated with sorafenib in a randomized, multicenter phase III trial.¹⁰ In this study, we aimed to analyze various potentially prognostic serum cytokines involved in cancer angiogenesis and cell proliferation using the multiplex immunoassay method before treatment and 4 weeks after axitinib initiation in patients with mRCC. We comprehensively explored biomarkers which can predict the clinical effect and prognosis in patients with mRCC treated with axitinib.

2 | MATERIAL AND METHODS

2.1 | Patients

From September 2012 to October 2015, 44 patients with mRCC at the Akita University Hospital were enrolled. An approval (#924) was

obtained by Akita University Hospital Institutional Review Board in accordance with the ethical standards based on the Declaration of Helsinki and its later amendments. Written informed consent was obtained by all the patients who participated in this study. Serum samples were obtained before treatment and 4 weeks after axitinib initiation. Patient characteristics are presented in Table 1. The International Metastatic Renal Cell Carcinoma Database Consortium (IMDC) risk classification at the axitinib initiation treatment was favorable in 11 (25.0%), intermediate in 30 (68.2%), and poor in 7 (15.9%). Twenty-six (59.1%) patients received no other therapies before axitinib. Axitinib treatment was initiated at 10 mg/day twice daily; thereafter, the dosage was increased or decreased according to the discretion of the attending physician based on serum axitinib level, adverse events, and treatment effect. Evaluation of the therapeutic effect was based on the Response Evaluation Criteria in Solid Tumors v1.1.

2.2 | Quantitative analysis of serum biomarkers

Serum samples were centrifuged at 3000 revolutions per min for 10 minutes, and stored at -80°C prior to analysis. Beads array analysis using the Bio-Plex Pro Cancer Biomarker assay kit1 and kit2 (Bio-Rad, Hercules, California) was performed to measure 34 cytokines and tumor growth factors.

Briefly, the capture antibody-coupled beads were first incubated with antigen standards, quality control samples, and serum samples in 96-well plates, followed by incubation with biotinylated detection antibodies. Samples were diluted 1:4 using sample diluent. After washing the unbound biotinylated detection antibodies, the beads were incubated with a reporter streptavidin-phycoerythrin (SA-PE) conjugate. Following the removal of excess SA-PE, the beads were passed through the 2-laser flow cytometer Bio-Plex array reader (Bio-Plex 200 system, Bio-Rad), which measures the fluorescence of the bead and the bound SA-PE. Details of the procedure have been described previously.¹¹ Assay incubations were performed at room temperature. All washes were performed using the Bio-Plex Pro wash station. Data acquisition was performed using Bio-Plex manager TM 6.0. Using the automatic calibration curve optimization function, the recovery rate was regressed to be in the range of approximately 70% to 130%. All samples were assayed in duplicate.

The following biomarkers were determined using the Bio-Plex Pro Human Cancer Biomarker Panel kit1 (#171-AC500M, Bio-

TABLE 1 Patients characteristics of the 44 patients with metastatic renal cell carcinoma treated with axitinib

		No. of patients (%) n = 44
Gender	Male	31 (70.5)
	Female	13 (29.5)
Age	Median [range]	66.5 [24-83]
BMI	Median [range]	22.7 [16.1-31.8]
IMDC risk group classification	Favorable	8 (18.2)
	Intermediate	24 (54.5)
	Poor	7 (15.9)
	Not available	5 (11.4)
Histological type	Clear cell	36 (81.8)
	Chromophobe	2 (4.5)
	Xp translocation	4 (9.1)
	Sarcomatoid	2 (4.5)
Nephrectomy	Yes	41 (93.2)
	No	3 (6.8)
Target organ	Lung	29 (65.9)
	Lymph node	14 (31.8)
	Bone	11 (25.0)
	Liver	5 (11.4)
Previous treatment	Yes	18 (40.9)
	At least one previous molecular-targeted agent	12 (66.7)
	Sunitinib	11 (61.1)
	Sorafenib	4 (22.2)
	Everolimus	7 (38.9)
	Temsirolimus	1 (5.6)
	Cytokines only	6 (33.3)
	No	26 (59.1)

Rad): soluble epidermal growth factor receptor (sEGFR), fibroblast growth factor basic (FGF-basic), soluble VEGF receptor (sVEGFR)-1, sVEGFR-2, platelet endothelial cell adhesion molecule-1 (PECAM-1), platelet-derived growth factor-AB/BB (PDGF-AB/BB), granulocyte-colony stimulating factor (G-CSF), hepatocyte growth factor (HGF), tyrosine kinase sHER-2/neu (erbB-2), tyrosine kinase sTIE2, sIL-6R α , *folistatin*, prolactin (PRL), leptin, and osteopontin. In addition, the following biomarkers were determined using the Bio-Plex Pro Human Cancer Biomarker Panel kit2 (#171-AC600M, Bio-Rad): VEGF-A, VEGF-C, VEGF-D, epidermal growth factor receptor (EGFR), heparin-binding epidermal growth factor-like growth factor (HB-EGF), *placental growth factor* (PLGF), *transforming growth factor- α* (TGF- α), tumor necrosis factor- α (TNF- α), insulin-like growth factor-binding protein 1 (IGFBP-1), soluble Fas ligand (sFASL), IL-6, IL-8, IL-18, plasminogen activator inhibitor-1 (PAI-1), urokinase plasminogen activator (uPA), angiotensin-2, sCD40L, and endoglin.

2.3 | Immunohistochemistry staining

Tumor specimens obtained by radical nephrectomy or biopsy were fixed in 20% formalin, embedded in paraffin, and evaluated for expression of PAI-1. Specimens were sliced into 3 μ m sections and immunohistochemically analyzed using anti-PAI-1 antibody (#66705, Abcam, Cambridge, UK). Peroxidase and 3,3-diaminobenzidine (DAB) were used as labeling enzyme and chromogenic substrate, respectively. Immunohistochemistry (IHC) staining was assessed using an automated quantitative pathology imaging system workstation (Mantra, PerkinElmer, Waltham, Massachusetts). DAB-positive cells were detected, and the staining intensity was scored using inForm ver. 2.3 software (PerkinElmer). Five representative areas were photographed with a 400-fold field of view, and nuclei were automatically recognized. Staining intensity was measured radially from the nucleus, and DAB staining was recognized around the cell membrane (Figure S1). The positive threshold for staining intensity per cell was defined as $\geq 25\%$ of the maximum staining intensity. The percentage of cells exceeding the threshold was counted, and the average value of the five visualized areas was scored as the final IHC staining intensity.

2.4 | Statistical analysis

The Kolmogorov-Smirnov test was used for nonparametric analysis of the serum biomarkers because of their nonnormal distribution. The relationships between serum biomarker level, treatment response, IHC staining intensity, and pathological parameters were evaluated using the Mann-Whitney *U* test. Bonferroni's correction was applied in the multiple comparison. Fisher's exact test was used to examine the proportion of patients between groups. The Kaplan-Meier method was used to plot time-to-event curves, and statistical significance was estimated using the log-rank test. The Cox proportional hazard model was used to determine independent prognostic factors of PFS and OS. $P < .05$ was considered as statistically significant. All statistical analyses were performed using SPSS statistics version 23 (IBM, New York).

3 | RESULTS

3.1 | Change in serum biomarker levels from pre-treatment to 4 weeks after axitinib initiation

Among the 34 measured cancer-related biomarkers, the median serum level of sTIE2, sVEGFR-1, sVEGFR-2, and Ang2 significantly decreased from pre-treatment to 4 weeks after axitinib initiation ($P < .001$, $P = .036$, $P < .001$, and $P = .006$, respectively; Table 2).

In contrast, the median serum level of sEGFR and PRL significantly increased from pre-treatment to 4 weeks after axitinib initiation ($P = .032$ and $P = .010$, respectively; Table 2). Using Bonferroni's correction, only sTIE2, sVEGFR-2, and PRL were significantly decreased or increased. The number of patients for each serum biomarker who exhibited a decrease or increase in level is shown in Table 2.

TABLE 2 List of the determined biomarkers and their serum level of pre-treatment and 4 weeks after initiation of axitinib

Protein name	Abbreviations	Pre-treatment		4 weeks after initiation of axitinib		P value	Number of patients for change in the serum level	
		Median (pg/mL)	Range	Median (pg/mL)	Range		Increased (n)	Decreased (n)
Bio-Plex Pro Human Cancer Biomarker Panel kit1								
Soluble epidermal growth factor receptor	sEGFR	14 779	12 669-18 295	15 200	13 386-20 398	.032	29	15
Fibroblast growth factor basic	FGF-basic	194	161-218	183	160-215	.090	17	27
Follistatin	Follistatin	707	506-948	629	497-1279	.375	23	21
Granulocyte-colony stimulating factor	G-CSF	82	60-93	76	62-90	.255	19	25
Tyrosine kinase soluble HER-2/neu	erbB-2	2186	1705-3348	2754	1618-3288	.273	24	20
Hepatocyte growth factor	HGF	1246	1022-2783	1305	1050-3174	.666	23	21
Soluble IL-6R α	sIL-6R α	10 180	8227-11 940	10 507	8329-12 732	.161	28	16
Leptin	Leptin	1907	1016-4364	2134	924-3545	.788	21	23
Osteopontin	OPN	70 999	45 785-90 563	69 869	47 384-94 053	.972	22	22
Platelet-derived growth factor-AB/BB	PDGF-AB/BB	2732	1941-4126	2796	1939-3815	.735	22	22
Platelet endothelial cell adhesion molecule-1	PECAM-1	2981	2539-4093	3257	2662-3849	.926	26	18
Prolactin	PRL	6029	4378-11 048	8036	5323-17 673	.010	33	11
Stem cell factor	SCF	219	197-267	219	193-247	.076	16	28
Tyrosine kinase soluble TIE2	sTIE-2	6168	5137-8635	5510	4082-7099	<.001	9	35
Soluble vascular endothelial growth factor receptor-1	sVEGFR-1	219	138-304	188	140-257	.036	17	27
Soluble vascular endothelial growth factor receptor-1	sVEGFR-2	3558	2728-4098	2830	2209-3217	<.001	7	37
Bio-Plex Pro Human Cancer Biomarker Panel kit2								
Angiopoietin-2	Ang2	954	567-1306	751	292-1366	.006	13	31
Soluble CD40 ligand	sCD40L	412	286-487	390	308-495	.797	22	22
Epidermal growth factor receptor	EGF	58	29-89	62	33-99	.161	28	16
Endoglin	ENG	906	459-1197	817	413-1186	.138	19	25
Soluble Fas ligand	sFASL	298	259-396	278	226-420	.118	15	29
Heparin-binding epidermal growth factor-like growth factor	HB-EGF	79	54-96	71	46-102	.197	21	23
Insulin-like growth factor-binding protein 1	IGFBP-1	12 372	4731-18 729	11 605	3447-28 333	.135	26	18
Interleukin-6	IL-6	80	33-102	68	26-108	.930	23	21
Interleukin-8	IL-8	24	13-29	24	12-34	.718	23	21
Interleukin-18	IL-18	135	105-182	160	91-207	.243	23	21
Plasminogen activator inhibitor-1	PAI-1	110 156	74 073-165 898	107 590	76 894-147 861	.991	24	20
Placental growth factor	PLGF	86	43-128	102	52-141	.067	30	14

TABLE 2 (Continued)

Protein name	Abbreviations	Pre-treatment		4 weeks after initiation of axitinib		P value	Number of patients for change in the serum level	
		Median (pg/mL)	Range	Median (pg/mL)	Range		Increased (n)	Decreased (n)
Transforming growth factor- α	TGF- α	60	46-81	52	38-86	.700	21	23
Tumor necrosis factor- α	TNF- α	44	16-67	39	14-61	.280	20	24
Urokinase plasminogen activator	uPA	228	74-340	210	69-371	.981	21	23
Soluble vascular endothelial growth factor A	VEGF-A	580	459-754	610	382-862	.401	25	19
Soluble vascular endothelial growth factor C	VEGF-C	959	671-1075	921	580-1167	.815	24	20
Soluble vascular endothelial growth factor D	VEGF-D	862	498-1633	753	466-1600	.155	19	25




FIGURE 1 Kaplan-Meier curves comparing, A, progression-free survival, and B, overall survival in patients with decreased or increased serum plasminogen activator inhibitor-1 (PAI-1) level from pre-treatment to 4 weeks after axitinib initiation

3.2 | Relationship between serum biomarker levels and treatment response

The treatment responses of 42 patients treated with axitinib were partial remission (PR) in 16 (38.1%) patients, stable disease (SD) in 20 (47.6%), and progressive disease (PD) in 6 (14.3%). Two patients were excluded because of unknown response. The median serum PDGF-AB/BB and sVEGFR-2 levels at baseline were significantly higher in the six patients with PD than in the 36 patients with PR or SD ($P = .040$ and $P = .003$, respectively); however, the baseline median

serum PAI-1 level was significantly lower in the patients with PD than those with PR or SD ($P = .048$) (Table S1). Using Bonferroni's correction, there was no significant relationship.

The proportion of patients with decreased serum level of PAI-1 and IL-18 from pre-treatment to 4 weeks after axitinib initiation was significantly higher in patients with PR or SD compared to those with PD ($P = .022$ and $P = .022$, respectively; Table S2). The proportion of patients with decreased serum levels of endoglin, IL-6, and VEGF-A from pre-treatment to 4 weeks after axitinib initiation was significantly higher in patients with PR than those with SD or PD ($P = .011$,

TABLE 3 Cox proportional hazard model to predict the shorter progression-free survival using baseline clinical parameter and change in the serum biomarker level from pre-treatment to 4 weeks after initiation of axitinib

Variable	Univariate analysis			Multivariate analysis (stepwise)		
	HR	95% CI	P value	HR	95%CI	P value
Age (<median vs >median)	0.747	0.346-1.611	.456			
Gender (male vs female)	1.048	0.441-2.493	.915			
BMI (<25 vs \geq 25)	0.788	0.359-1.730	.553			
Previous treatment (no vs yes)	0.850	0.349-1.831	.678			
pT (\geq pT2 vs pT1)	1.508	0.627-3.628	.359			
cN (\geq cN1 vs cN0)	5.476	2.039-14.704	.001	10.616	3.287-34.280	<.001
LVI (yes vs no)	1.226	0.409-3.672	.716			
Grade (G2-3 vs G1)	1.141	0.586-2.219	.699			
Number of metastasis (\geq 3 vs 0-2)	1.937	0.838-4.477	.122			
Lung metastasis (yes vs no)	1.019	0.441-2.353	.965			
Liver metastasis (yes vs no)	3.236	1.180-8.875	.022	2.854	0.843-9.662	.092
Bone metastasis (yes vs no)	1.890	0.823-4.338	.133			
CRP (\geq ULN vs <ULN)	1.114	0.486-2.554	.798			
Alb (<LLN vs >LLN)	2.630	0.991-6.981	.052			
Hb (<LLN vs >LLN)	1.859	0.858-4.028	.112			
Thrombocyte(<ULN vs \geq ULN)	1.802	0.674-4.819	.241			
sEGFR (increased vs decreased)	0.787	0.348-1.780	.565			
FGF-basic (increased vs decreased)	1.217	0.686-2.158	.501			
Follistatin (increased vs decreased)	0.859	0.396-1.863	.700			
G-CSF (increased vs decreased)	1.124	0.525-2.406	.763			
erbB-2 (increased vs decreased)	1.039	0.471-2.291	.925			
HGF (increased vs decreased)	1.492	0.689-3.230	.310			
IL-6R α (increased vs decreased)	1.573	0.687-3.605	.284			
Leptin (increased vs decreased)	0.953	0.446-2.036	.900			
OPN (increased vs decreased)	1.078	0.503-2.313	.847			
PDGF-AB/BB (increased vs decreased)	0.860	0.402-1.837	.697			
PECAM-1 (increased vs decreased)	1.377	0.611-3.104	.441			
PRL (increased vs decreased)	1.233	0.519-2.929	.635			
SCF(increased vs decreased)	1.002	0.458-2.193	.996			
TIE2 (increased vs decreased)	0.711	0.283-1.782	.466			
sVEGFR-1 (increased vs decreased)	0.764	0.378-1.541	.451			
sVEGFR-2 (increased vs decreased)	0.839	0.313-2.245	.726			
Ang2 (increased vs decreased)	0.809	0.341-1.921	.631			
sCD40L (increased vs decreased)	2.135	0.956-4.770	.064			
EGF (increased vs decreased)	1.809	0.763-4.289	.178			
ENG (increased vs decreased)	1.667	0.780-3.563	.188			
sFASL (increased vs decreased)	1.457	0.665-3.193	.347			
HB-EGF (increased vs decreased)	2.233	1.027-4.854	.043	1.937	0.208-60.373	.561
IGFBP-1 (increased vs decreased)	1.359	0.619-2.986	.444			
IL-6 (increased vs decreased)	2.328	1.053-5.143	.037	1.037	0.332-3.237	.949
IL-8 (increased vs decreased)	1.935	0.879-4.258	.101			
IL-18 (increased vs decreased)	1.675	0.759-3.694	.201			
PAI-1 (increased vs decreased)	2.412	1.075-5.412	.027	3.896	1.306-11.623	.015
PLGF (increased vs decreased)	2.671	1.008-7.075	.048	2.018	0.547-8.127	.279

TABLE 3 (Continued)

Variable	Univariate analysis			Multivariate analysis (stepwise)		
	HR	95% CI	P value	HR	95%CI	P value
TGF- α (increased vs decreased)	2.485	1.114-5.546	.026	0.912	0.089-9.039	.938
TNF- α (increased vs decreased)	1.995	0.928-4.291	.077			
uPA (increased vs decreased)	1.444	0.693-3.008	.327			
VEGF-A (increased vs decreased)	1.435	0.656-3.142	.366			
VEGF-C (increased vs decreased)	1.924	0.875-4.233	.104			
VEGF-D (increased vs decreased)	1.608	0.753-3.432	.220			

TABLE 4 Cox proportional hazard model to predict the shorter overall survival using baseline clinical parameter and change in the serum biomarker level from pre-treatment to 4 weeks after initiation of axitinib

Variable	Univariate			Multivariate		
	HR	95% CI	P value	HR	95%CI	P value
Age (<median vs >median)	0.480	0.174-1.324	.156			
Gender (male vs female)	0.854	0.274-2.658	.785			
BMI (<25 vs \geq 25)	0.602	0.208-1.745	.350			
Previous treatment (no vs yes)	0.534	0.182-1.568	.253			
pT (\geq pT2 vs pT1)	1.233	0.386-3.942	.724			
cN (\geq cN1 vs cN0)	4.691	1.562-14.089	.006	2.292	0.483-10.883	.297
LVI (yes vs no)	1.494	0.326-6.853	.606			
Grade (G2-3 vs G1)	1.439	0.597-3.473	.418			
Number of metastasis (\geq 3 vs 0-2)	4.104	1.487-11.321	.006	2.709	0.357-20.533	.335
Lung metastasis (yes vs no)	0.912	0.311-2.674	.867			
Liver metastasis (yes vs no)	2.841	0.904-8.924	.074			
Bone metastasis (yes vs no)	3.255	1.198-8.846	.021	2.472	0.370-16.492	.35
CRP (\geq ULN vs <ULN)	3.102	0.703-13.684	.135			
Alb (<LLN vs >LLN)	3.417	0.769-15.175	.106			
Hb (<LLN vs >LLN)	3.382	1.090-10.496	.035	1.996	0.534-7.453	.304
Thrombocyte(<ULN vs \geq ULN)	3.046	0.957-9.699	.059			
sEGFR (increased vs decreased)	0.753	0.273-2.079	.584			
FGF-basic (increased vs decreased)	1.119	0.508-2.464	.781			
Follistatin (increased vs decreased)	0.969	0.363-2.586	.949			
G-CSF (increased vs decreased)	0.622	0.215-1.799	.381			
erbB-2 (increased vs decreased)	0.701	0.261-1.880	.481			
HGF (increased vs decreased)	1.753	0.637-4.824	.277			
IL-6R α (increased vs decreased)	2.130	0.684-6.637	.192			
Leptin (increased vs decreased)	1.203	0.451-3.210	.712			
OPN (increased vs decreased)	1.498	0.533-4.212	.443			
PDGF-AB/BB (increased vs decreased)	0.678	0.245-1.874	.454			
PECAM-1 (increased vs decreased)	0.906	0.336-2.443	.846			
PRL (increased vs decreased)	0.846	0.294-2.437	.757			
SCF(increased vs decreased)	1.729	0.647-4.625	.275			
TIE2 (increased vs decreased)	0.651	0.185-2.289	.503			
sVEGFR-1 (increased vs decreased)	0.634	0.244-1.647	.349			
sVEGFR-2 (increased vs decreased)	1.104	0.286-3.598	.983			
Ang2 (increased vs decreased)	1.279	0.455-3.595	.641			

(Continues)

TABLE 4 (Continued)

Variable	Univariate			Multivariate		
	HR	95% CI	P value	HR	95%CI	P value
sCD40L (increased vs decreased)	1.173	0.434-3.173	.753			
EGF (increased vs decreased)	0.804	0.292-2.219	.674			
ENG (increased vs decreased)	1.175	0.441-3.133	.747			
sFASL (increased vs decreased)	1.228	0.443-3.399	.693			
HB-EGF (increased vs decreased)	1.486	0.549-4.025	.436			
IGFBP-1 (increased vs decreased)	1.237	0.449-3.408	.680			
IL-6 (increased vs decreased)	2.349	0.813-6.783	.115			
IL-8 (increased vs decreased)	0.916	0.331-2.531	.865			
IL-18 (increased vs decreased)	1.539	0.559-4.240	.404			
PAI-1 (increased vs decreased)	3.376	1.086-10.497	.036	5.316	1.154-24.488	.032
PLGF (increased vs decreased)	1.424	0.453-4.474	.545			
TGF- α (increased vs decreased)	1.486	0.549-4.025	.436			
TNF- α (increased vs decreased)	1.130	0.424-3.015	.807			
uPA (increased vs decreased)	2.240	0.819-6.123	.116			
VEGF-A (increased vs decreased)	1.057	0.383-2.918	.915			
VEGF-C (increased vs decreased)	1.508	0.547-4.152	.427			
VEGF-D (increased vs decreased)	0.846	0.312-2.298	.743			

$P = .025$, and $P = .029$, respectively; Table S2). Using Bonferroni's correction, there was no significant relationship.

3.3 | Relationship between serum biomarker levels and PFS and OS

The presence of lymph node swelling on initial imaging studies (cN1) and baseline serum leptin level lower than the median were independent factors related to worse PFS in multivariate analysis ($P < .001$ and $P = .026$; Table S3). No independent factor related to OS was found using baseline serum biomarker level (Table S4).

Patients with increased serum PAI-1 level from pre-treatment to 4 weeks after axitinib initiation had significantly shorter PFS and OS than those with decreased serum PAI-1 (15.0 months vs 5.1 months, $P = .027$ and 34.9 months vs 14.2 months, $P = .026$, respectively; Figure 1A,B). The presence of lymph node swelling on initial imaging studies (cN1) and increased serum PAI-1 level from pre-treatment to 4 weeks after axitinib initiation were independent prognostic factors for shorter PFS ($P < .001$ and $P = .015$, respectively; Table 3). Increased serum PAI-1 level from pre-treatment to 4 weeks after axitinib initiation was also an independent prognostic marker for shorter OS ($P = .032$; Table 4).

3.4 | Relationship between IHC staining intensity and clinical parameters

Of the 44 patients enrolled in this study, 41 (93.2%) underwent radical nephrectomy and 3 (6.8%) underwent tumor biopsy. IHC analysis

using PAI-1 antibody was available in 39 specimens from 36 nephrectomies and 3 biopsies. The median IHC staining intensity of PAI-1 was significantly higher in patients with metastatic disease at the time of diagnosis than those with nonmetastatic disease ($P = .010$; Table 5), as well as in patients with Fuhrman grade ≥ 3 tumors than in those with grade ≤ 2 ($P = .026$; Table 5). There was no significant relationship between PAI-1 staining intensity and PFS or OS (Figure S2), and between PAI-1 staining intensity and serum baseline PAI-1 level ($r^2 = 0.053$, $\rho = -0.02$, $P = .904$).

4 | DISCUSSION

The multiplex immunoassay method is a beads array in which various antibodies are loaded on the beads measured by flow cytometry. Previous reports have comprehensively measured angiogenic factors using serum samples from patients with colorectal, ovarian and small cell lung cancer¹²⁻¹⁴ and urine samples from patients with bladder cancer.^{15,16} However, few studies have explored biomarkers as predictive factors in patients with metastatic disease using multiplex immunoassay techniques. Although we expected biomarkers other than sVEGFRs to show predictive value in this study, serum PAI-1 level was the only biomarker associated with therapeutic effect, PFS, and OS after axitinib treatment in patients with mRCC.

PAI-1 usually exists in vascular endothelial cells, liver, platelets, and adipocytes, and functions as the principal inhibitor of urokinase-type plasminogen activator (uPA) and its receptor (uPAR) system in fibrinolysis. Furthermore, $\geq 90\%$ of PAI-1 is contained in platelets and released into the bloodstream under conditions of vascular endothelial

biomarker level or change in level after treatment. These results are partially consistent with previous studies that reported sVEGFR-2 and sVEGFR-3 levels were significant prognostic factors after sunitinib treatment in patients with mRCC.^{6,7} Although serum PAI-1 and sVEGFRs have been identified as markers of tumor hypoxia, and might be affected by systemic VEGF-directed inhibitors,^{28,32} serum PAI-1 level may be a more useful prognostic biomarker than serum sVEGFRs in this axitinib study.

There are several important limitations of this study. First, PAI-1 is ideally measured in plasma, however we used serum samples in this study, which might affect the results. Second, the PAI-1 level measured in this study was not pure PAI-1 but a complex in the blood. The antibody on the beads of the Bio-Plex Pro Human Cancer Biomarker Panel 2 in this study is an anti-total PAI-1 antibody, which measures the sum of the active type, latent type, vitronectin complex, tissue-type plasminogen activator complex, and uPA complex. Third, 40% of patients received multiple therapies prior to axitinib treatment, which might affect the interpretation of the results. To verify our results, future studies measuring plasma PAI-1 level in larger RCC cohorts should be conducted.

5 | CONCLUSIONS

The initial changes in serum PAI-1 level at the early stage of axitinib treatment could be a useful prognostic biomarker in patients with mRCC.

ACKNOWLEDGMENT

The authors thank Yoko Mitobe, Yukiko Sugiyama, and Yuka Izumida for their contribution of clinical sample collections and preparations.

FUNDING

This work was supported by the grant numbers 25293332, 16H02679, and 23590168 from the Japanese Society for the Promotion of Science and AMED-CREST, Japan Agency for Medical Research and Development (AMED).

CONFLICT OF INTEREST

The authors declare no conflicts of interest.

AUTHOR CONTRIBUTIONS

Conceptualization: Naoko Honma, Takamitsu Inoue, and Norihiko Tsuchiya

Data curation: Naoko Honma, Takamitsu Inoue, and Norihiko Tsuchiya

Formal analysis: Naoko Honma and Takamitsu Inoue

Funding acquisition: Takamitsu Inoue, Norihiko Tsuchiya, and Tomonori Habuchi

Investigation: Naoko Honma, Mingguo Huang, and Takamitsu Inoue

Methodology: Naoko Honma, Takamitsu Inoue, and Norihiko Tsuchiya

Project administration: Naoko Honma and Takamitsu Inoue

Resources: Naoko Honma, Takamitsu Inoue, Atsushi Koizumi, Ryohei Yamamoto, Taketoshi Nara, Sohei Kanda, Mitsuru Saito, Kazuyuki Numakura, and Shintaro Narita

Software: Naoko Honma, Mingguo Huang, and Takamitsu Inoue

Supervision: Shintaro Narita, Shigeru Satoh, and Tomonori Habuchi

Validation: Shintaro Narita

Visualization: Naoko Honma and Takamitsu Inoue

Writing - original draft preparation: Naoko Honma

Writing - review and editing: Takamitsu Inoue, Shintaro Narita, Shigeru Satoh, and Tomonori Habuchi

All authors have read and approved the final version of the manuscript.

TRANSPARENCY STATEMENT

The corresponding author, Takamitsu Inoue, affirms that this manuscript is an honest, accurate, and transparent account of the study being reported; that no important aspects of the study have been omitted; and that any discrepancies from the study as planned (and, if relevant, registered) have been explained.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are openly available in "figshare" at <https://figshare.com/s/ea7a0931565d9b36f1e2>, DOI: 10.6084/m9.figshare.12049560.

ORCID

Takamitsu Inoue  <https://orcid.org/0000-0002-1362-4344>

Kazuyuki Numakura  <https://orcid.org/0000-0002-5899-1738>

REFERENCES

1. Tsushima T. Epidemiology of renal cell carcinoma in Japan. *Nippon Rinsho*. 2017;75:23-26.
2. Sun M, Thuret R, Abdollah F, et al. Age-adjusted incidence, mortality, and survival rates of stage-specific renal cell carcinoma in North America: a trend analysis. *Eur Urol*. 2011;59:135-141.
3. Lalani AA, McGregor BA, Albiges L, et al. Systemic treatment of metastatic clear cell renal cell carcinoma in 2018: current paradigms, use of immunotherapy, and future directions. *Eur Urol*. 2019;75:100-110.
4. Wahlgren T, Harmenberg U, Sandstrom P, et al. Treatment and overall survival in renal cell carcinoma: a Swedish population-based study (2000-2008). *Br J Cancer*. 2013;108:1541-1549.
5. Igarashi R, Inoue T, Fujiyama N, et al. Contribution of UGT1A1 genetic polymorphisms related to axitinib pharmacokinetics to safety and efficacy in patients with renal cell carcinoma. *Med Oncol*. 2018; 35:51.
6. Deprimo SE, Bello CL, Smeraglia J, et al. Circulating protein biomarkers of pharmacodynamic activity of sunitinib in patients with metastatic renal cell carcinoma: modulation of VEGF and VEGF-related proteins. *J Transl Med*. 2007;5:32.
7. Rini BI, Michaelson MD, Rosenberg JE, et al. Antitumor activity and biomarker analysis of sunitinib in patients with bevacizumab-refractory metastatic renal cell carcinoma. *J Clin Oncol*. 2008;26(22): 3743-3748.
8. Murukesh N, Dive C, Jayson GC. Biomarkers of angiogenesis and their role in the development of VEGF inhibitors. *Br J Cancer*. 2010; 102:8-18.

9. Gruenewald V, Beutel G, Schuch-Jantsch S, et al. Circulating endothelial cells are an early predictor in renal cell carcinoma for tumor response to sunitinib. *BMC Cancer*. 2010;10:69.
10. Motzer RJ, Escudier B, Tomczak P, et al. Axitinib versus sorafenib as second-line treatment for advanced renal cell carcinoma: overall survival analysis and updated results from a randomized phase 3 trial. *Lancet Oncol*. 2013;14:552-562.
11. Li D, Chiu H, Gupta V, Chan DW. Validation of a multiplex immunoassay for serum angiogenic factors as biomarkers for aggressive prostate cancer. *Clin Chim Acta*. 2012;413:1506-1511.
12. Villar-Vázquez R, Padilla G, Fernández-Aceñero MJ, et al. Development of a novel multiplex beads-based assay for autoantibody detection for colorectal cancer diagnosis. *Proteomics*. 2016;16:1280-1290.
13. Horala A, Swiatly A, Matysiak J, et al. Diagnostic value of serum angiogenesis markers in ovarian cancer using multiplex immunoassay. *Int J Mol Sci*. 2017;18:123.
14. Klupczynska A, Dereziński P, Matysiak J, et al. Determination of 16 serum angiogenic factors in stage I non-small cell lung cancer using a bead-based multiplex immunoassay. *Biomed Pharmacother*. 2017;88:1031-1037.
15. Shimizu Y, Furuya H, Greenwood PB, et al. A multiplex immunoassay for the non-invasive detection of bladder cancer. *J Transl Med*. 2016;14:31.
16. Goodison S, Ogawa O, Matsui Y, et al. A multiplex urinary immunoassay for bladder cancer detection: analysis of a Japanese cohort. *J Transl Med*. 2016;14:287.
17. Tjärnlund-Wolf A, Brogren H, Lo EH, Wang X. Plasminogen activator inhibitor-1 and thrombotic cerebrovascular diseases. *Stroke*. 2012;43(10):2833-2839.
18. Dass K, Ahmad A, Azmi AS, Sarkar SH, Sarkar FH. Evolving role of uPA/uPAR system in human cancers. *Cancer Treat Rev*. 2008;34:122-136.
19. Carter JC, Church FC. Obesity and breast cancer: the roles of peroxisome proliferator-activated receptor- γ and plasminogen activator inhibitor-1. *PPAR Res*. 2009;2009:1-13. <https://doi.org/10.1155/2009/345320>.
20. Hofmann R, Lehmer A, Buresch M, Hartung R, Ulm K. Clinical relevance of urokinase plasminogen activator, its receptor, and its inhibitor in patients with renal cell carcinoma. *Cancer*. 1996;78:487-492.
21. Hofmann R, Lehmer A, Hartung R, Robrecht C, Buresch M, Grothe F. Prognostic value of urokinase plasminogen activator and plasminogen activator inhibitor-1 in renal cell cancer. *J Urol*. 1996;155:858-862.
22. Ohba K, Miyata Y, Kanda S, Koga S, Hayashi T, Kanetake H. Expression of urokinase-type plasminogen activator, urokinase-type plasminogen activator receptor and plasminogen activator inhibitors in patients with renal cell carcinoma: correlation with tumor-associated macrophage and prognosis. *J Urol*. 2005;174:461-465.
23. Zubac DP, Wentzel-Larsen T, Seidal T, Bostad L. Type 1 plasminogen activator inhibitor (PAI-1) in clear cell renal cell carcinoma (CCRCC) and its impact on angiogenesis, progression and patient survival after radical nephrectomy. *BMC Urol*. 2010;10:20.
24. Choi JW, Lee JH, Park HS, Kim YS. PAI-1 expression and its regulation by promoter 4G/5G polymorphism in clear cell renal cell carcinoma. *J Clin Pathol*. 2011;64:893-897.
25. Chautard D, Dalifard I, Chassevent A, et al. Prognostic value of uPA, PAI-1, and DNA content in adult renal cell carcinoma. *Urology*. 2004;63:1055-1060.
26. Swiercz R, Wolfe JD, Zaher A, Jankun J. Expression of the plasminogen activation system in kidney cancer correlates with its aggressive phenotype. *Clin Cancer Res*. 1998;4:869-877.
27. Fuessel S, Erdmann K, Taubert H, et al. Prognostic impact of urokinase-type plasminogen activator system components in clear cell renal cell carcinoma patients without distant metastasis. *BMC Cancer*. 2014;14:974.
28. Lara PN Jr, Quinn DI, Margolin K, et al. California cancer consortium. SU5416 plus interferon alpha in advanced renal cell carcinoma: a phase II California cancer consortium study with biological and imaging correlates of angiogenesis inhibition. *Clin Cancer Res*. 2003;9(13):4772-4781.
29. Ferroni P, Roselli M, Portarena I, et al. Plasma plasminogen activator inhibitor-1 (PAI-1) levels in breast cancer - relationship with clinical outcome. *Anticancer Res*. 2014;34:1153-1161.
30. Grebenchtchikov N, Maguire TM, Riisbro R, et al. Measurement of plasminogen activator system components in plasma and tumor tissue extracts obtained from patients with breast cancer: an EORTC receptor and biomarker group collaboration. *Oncol Rep*. 2005;14:235-239.
31. Masuda T, Hattori N, Senoo T, et al. SK-216, an inhibitor of plasminogen activator inhibitor-1, limits tumor progression and angiogenesis. *Mol Cancer Ther*. 2013;12(11):2378-2388.
32. Koong AC, Denko NC, Hudson KM, et al. Candidate genes for the hypoxic tumor phenotype. *Cancer Res*. 2000;60(4):883-887.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of this article.

How to cite this article: Honma N, Inoue T, Tsuchiya N, et al. Prognostic value of plasminogen activator inhibitor-1 in biomarker exploration using multiplex immunoassay in patients with metastatic renal cell carcinoma treated with axitinib. *Health Sci Rep*. 2020;3:e197. <https://doi.org/10.1002/hsr2.197>